



US007452696B2

(12) **United States Patent**
Chen et al.

(10) **Patent No.:** **US 7,452,696 B2**
(45) **Date of Patent:** **Nov. 18, 2008**

(54) **RECOMBINANT PLASMID AND METHOD FOR EXPRESSING HEPATITIS B VIRAL ANTIGENS AND VIRIONS IN VIVO**

2005/0197313 A1* 9/2005 Roelvink et al. 514/44

(75) Inventors: **Pei-Jer Chen**, Taipei (TW); **Li-Rung Huang**, Taipei (TW); **Hui-Lin Wu**, Taipei (TW); **Ding-Shinn Chen**, Taipei (TW)

(73) Assignee: **National Taiwan University**, Taipei (TW)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/553,975**

(22) Filed: **Oct. 27, 2006**

(65) **Prior Publication Data**

US 2008/0124364 A1 May 29, 2008

(51) **Int. Cl.**
C12N 15/09 (2006.01)
C12P 21/04 (2006.01)
A01N 63/00 (2006.01)
A61K 39/12 (2006.01)

(52) **U.S. Cl.** **435/70.1**; 424/204.1; 424/93.1; 435/69.3

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2004/0214329 A1* 10/2004 Kay et al. 435/455

OTHER PUBLICATIONS

Arad et al., Liver-Targeted Gene Therapy by SV40-Based Vectors Using the Hydrodynamic Injection Method, 2005, Human Gene Therapy, vol. 16, pp. 361-371.*

Priscilla I. Yang, Alana Althage, Josan Chung, Francis V. Chisari, Hydrodynamic Injection of Viral DNA: A Mouse Model of Acute Hepatitis B Virus Infection, PNAS, Oct. 15, 2002, 13825-13830, vol. 99, No. 21.

F. Liu, Y.K. Song, D. Liu; Hydrodynamics-Based Transfection in Animals by Systemic Administration of Plasmid DNA, Gene Therapy (1999) 6, 1258-1266.

* cited by examiner

Primary Examiner—Bruce Campell

Assistant Examiner—Benjamin P Blumel

(74) *Attorney, Agent, or Firm*—WPAT, P.C.; Anthony King

(57) **ABSTRACT**

Disclosed is a recombinant plasmid for expressing hepatitis B viral antigens in vivo, comprising an adeno-associated virus (AVV) vector and a replication-competent hepatitis B virus genome fragment. Mice hydrodynamically injected with the recombinant plasmid of the present invention show persistent expression of hepatitis B viral antigens for more than 6 months in the hepatocytes, thus a immuno-competent mouse model for persistent expression of hepatitis B antigens and also for human chronic hepatitis B virus infection is established, which can be applied in evaluation and elucidation of mechanism of chronic hepatitis and anti-viral drug discovery research.

8 Claims, 5 Drawing Sheets




Figure 1




Figure 2




Figure 3






Figure 4

(A)



(B)




Figure 5

**RECOMBINANT PLASMID AND METHOD
FOR EXPRESSING HEPATITIS B VIRAL
ANTIGENS AND VIRIONS IN VIVO**

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a method for expressing hepatitis B viral antigens and virions in vivo, especially relates to a recombinant plasmid for expressing all of the hepatitis B viral antigens and a method to construct an immunocompetent mouse model for persistently expressing hepatitis B viral antigens by using the recombinant plasmid.

2. The Prior Arts

It is estimated that 350 million people are chronic carriers of hepatitis B virus worldwide, and there are approximately 1 million deaths each year caused by hepatitis B related liver diseases. Evidently, hepatitis B virus has tremendous impact on human health, therefore research for best treatment and new drug discovery enthusiastically become an important healthcare issue.

Hepatitis B virus (HBV) is a partially double-stranded DNA virus that infects human hepatocytes. Upon infection of host individual, the hepatitis B virus will first form a covalently closed circular DNA (cccDNA) within the cell, and then followed by a complicated viral replication process. Host individual infected by hepatitis B virus normally develops acute or chronic necroinflammatory of liver diseases, which consequently may transform into hepatocellular carcinoma. It is generally believed that HBV is not directly cytopathic to the hepatocytes, but HBV mediates the immune response of the infected hepatocytes and triggers the infected cells to express viral antigen, which resulting in host immune system attacking its own infected hepatocytes and eventually leading to cell apoptosis.

It is necessary and important to develop an appropriate animal model as a tool in search for new therapy and drug discovery, thus HBV transgenic mice have been proposed to serve as animal models in search for treatment of hepatitis and new drug discovery. The whole-genome HBV transgenic mice are created by delivery of a replication-competent HBV genome into mouse chromosome, in which the inserted viral genome is stably and persistently expressed in the liver tissue. However, it has been found that the mechanism of liver inflammation derived from such HBV transgenic mice is different from scenario of human hepatitis. Furthermore, such HBV transgenic mice are HBV antigen tolerant, thus do not exhibit normal immune response against HBV. Hence, it is concluded that such animal models are not ideal in study of HBV tolerance and clearance.

Chisari et al. devise an approach: by injecting hydrodynamically a recombinant plasmid containing the HBV whole genome into immunologically competent mice, to stimulate HBV replication and to induce hepatitis resembling infection process in human. It is proposed that construction of such readily manipulable animal models may be used for studies of HBV induced immune response and hepatitis. (Chisari et al., (2002) Proc. Natl. Acad. Sci. USA 99 (21), 13825-30). In their study, an inbred mouse B10.D2 that shows strong immune response to the HBV surface antigen is selected as a recipient, and pT-MCS-HBV1.3 plasmid is used as a vector to carry HBV gene. Chisari et al. successfully introduce the recombinant plasmid into mice and elicit immune response that resembles human hepatitis after HBV infection. However, there are disadvantages to this animal model, that is, short-term acute liver inflammatory response and clearance of immune system effect 14 days after transfection. These

constraints limit application of such animal model in studies such as long term monitoring of immune response after HBV infection and drug evaluation in chronic hepatitis.

SUMMARY OF THE INVENTION

In order to improve the shortcomings of currently existing acute hepatitis mouse model developed by Chisari et al. for HBV infection studies, it is necessary to develop a mouse model that can persistently express HBV antigens for studies such as hepatitis treatment and drug evaluation. Therefore, the primary object of the present invention is to provide a recombinant plasmid for persistently expressing HBV antigens in an immuno-competent host, which comprises an adeno-associated virus (abbreviated as AAV) vector, and a replication-competent genome fragment of HBV.

Another object of the present invention is to provide a method for expressing HBV antigen in vivo, which comprises the steps of:

- (1) providing a recombinant plasmid comprising an AVV vector, and a replication-competent HBV genome fragment; and
- (2) delivering the recombinant plasmid to a mouse and transfecting the hepatocytes of the mouse.

When the recombinant plasmid of the present invention is delivered into a mouse and transfected hepatocytes of the injected mouse, it can exist in an episomal form in the hepatocytes of the mouse for more than 200 days, and can be used as a HBV transcription template for production of viral antigens, replication intermediates, and mature virions which are released in bloodstream of the injected mouse. Because the characteristics of episomal form plasmid and the cccDNA of natural infected HBV are very similar, therefore, the recombinant plasmid of the present invention can be used for drug screening and evaluation of effectiveness of cccDNA clearance.

Furthermore, liver histological and serological status analysis of the carrier mouse receiving hydrodynamic injection of the recombinant plasmid of the present invention shows similar pattern to that of healthy HBV carriers in the tolerance stage. The results indicate that with this animal model, it is now possible to address some questions that are hardly studied in the conventional HBV transgenic mouse model, e.g. the tolerance development after HBV exposure, mechanisms of HBV acute exacerbation and the HBV cccDNA clearance in response to antiviral or immuno-modulator therapy etc. Furthermore, the effect of many HBV variants in regulating HBV replication or in contributing to viral persistence may also be conveniently studied in this model.

The present invention is further explained in the following embodiment illustration and examples. Those below should not, however, be considered to limit the scope of the invention, it is contemplated that modifications will readily occur to those skilled in the art, which modifications will be within the spirit of the invention and the scope of the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1(A) shows temporal profile of HBsAg titer in sera of C57BL/6 and BALB/c mice receiving hydrodynamic injection of the recombinant plasmid pAAV/HBV (1400). S/N ratio greater than 10 is defined as HBsAg-positive; ○:BALB/c mice; ●:C57BL/6 mice. FIG. 1(B) shows clearance rate of serum HBsAg of C57BL/6 mice receiving hydrodynamic injection of recombinant plasmid pAAV/HBV (1400) and

pGEM4Z/HBV (1400). S/N ratio greater than 10 is defined as HBsAg-positive. ○:pGEM4Z/HBV (1400) (n=8); ●:pAAV/ HBV (1400) (n=12).

FIG. 2 shows gel electrophoresis analysis of HBV replication intermediates and transcripts in the livers of C57BL/6 mice receiving hydrodynamic injection of plasmid pAAV/ HBV (1400). mGAPDH is the internal control of Northern hybridization, lane a is HBsAg-positive C57BL/6 mice, lane b is HBsAg-negative C57BL/6 mice which cleared HBV antigens after hydrodynamic injection of pAAV/ HBV (1400).

FIG. 3 shows immunohistochemical staining of the liver tissue of the mice receiving hydrodynamic injection of pAAV/ HBV (1400) at day 381. Experimental group is HBsAg-positive mice, and control group is HBsAg-negative mice, which cleared HBV antigens after hydrodynamic injection of pAAV/ HBV (1400). (A) HBsAg staining of experimental group, arrow indicates HBsAg-positive hepatocytes. (B) HBsAg staining of control group. (C) HBcAg staining of experimental group, arrow indicates HBcAg-positive hepatocytes. (D) HBcAg staining of control group.

FIG. 4 shows effect of age on HBsAg level of mice receiving hydrodynamic injection of pAAV/ HBV (1400). (A) Change of HBsAg titer in bloodstream; Δ:13-week old mice (n=9); ●:6.5-week old mice (n=7). (B) Ratio of HBsAg-positive mice (S/N ratio greater than 10 is defined as HBsAg-positive); Δ:13-week old mice (n=9); ●:6.5-week old mice (n=7).

FIG. 5 shows effect of gender on HBsAg level of C57BL/6 mice receiving hydrodynamic injection of pAAV/ HBV (1400). (A) Change of HBsAg titer in bloodstream; Δ:female mice (n=16); ●:male mice (n=12). (B) Percentage of HBsAg-positive mice (S/N ratio greater than 10 is defined as HBsAg-positive); Δ:female mice (n=16); ●:male mice (n=12).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention provides a recombinant plasmid for persistently expressing HBV antigens in vivo, which comprises an adeno-associated virus (AAV) vector and a replication-competent HBV genome fragment.

The abovementioned adeno-associated vector can be any AAV vector that is commercially available, and such commercial vector can be applied in the present invention as vector carrying HBV genome regardless its source. There is no limitation to the HBV genome fragment applied in the present invention, as long as the genome fragment can be inserted into the adeno-associated virus with any conventional method and can subsequently express or replicates, such as the genes encoding envelope proteins, core/precursor proteins, x protein or polymerase protein of HBV.

In one embodiment, the abovementioned HBV genome fragment is inserted in and flanked by inverted terminal repeats (abbreviated as ITRs) of AAV-GFP plasmid, to replace the Green fluorescent protein (GFP) gene fragment already existed in the plasmid. Thus constructed recombinant plasmid is designated as pAAV/ HBV (1400), which DNA sequence is listed as SEQ ID NO: 1. The recombinant plasmid of the present invention is deposited in the Food Industry Research and Development Institute (Taiwan, R.O.C.), with deposit ID No.: BCRC940490.

To persistently express HBV antigens in vivo using the abovementioned expression system, the method can begin with preparation of a recombinant plasmid (for example, pAAV/ HBV (1400) plasmid) and follow by delivering the

recombinant plasmid into mouse using any known skills in the art, and consequently the hepatocytes of the injected mouse is transfected.

Mice that are well known to those skilled in the art and with immune response level lower than inbred mice B10.D2 (H-2d haplotype) can be used in the present invention, such as C57BL/6 mice (H-2b haplotype), but it should not, however, be considered to limit the scope of the invention. Furthermore, the preferred haplotype of the receiving mouse is H-2^b.

In terms of delivery method of the recombinant plasmid, any approach that is well known in the arts of plasmid delivery and transfection of the liver cells can be applied in the present invention, but should not be considered to limit the scope of the present invention. In the present invention, for example, known skills such as hydrodynamic injection can be one of the methods for plasmid delivery. More specifically, in the example of the present invention, transfection of mouse liver cells with the recombinant plasmid is accomplished by hydrodynamic injection of the recombinant plasmid into the tail vein of mice. In order to allow the recombinant plasmid easily injected into tail vein of mice, the plasmid abovementioned is prepared in a biocompatible and non-immunogenic solution, such as phosphate buffer solution, but it should not be considered to limit the scope of the present invention.

Once the recombinant plasmid of the present invention is properly delivered into mice and the hepatocytes are transfected, the plasmids in the hepatocyte are existed as episomal form and are persistently served as templates for transcription for more than 200 days. Moreover the expression of HBV antigens and replication intermediates are also persistently expressed and detected for more than 26 weeks.

Furthermore, because the characteristics of episomal form of the present invention recombinant plasmid are similar to that of the cccDNA in natural HBV infection, liver serological and histological results of the carrier mice of the present invention resemble to that of healthy human carrier. Hence, the recombinant plasmid of the present invention and its application can be applied in construction of mouse models for persistent expression of HBV antigen for more than 6 months.

According to the previous studies of natural HBV infection in human, more than 90% acutely infected adults resolve all clinical symptoms, develop HBeAg- and HBsAg specific antibodies, clear HBeAg and HBsAg from the circulation and maintain lifelong protective immunity. In contrast to HBV infection during adulthood, perinatal HBV infection typically results in chronic hepatitis. In the present invention, different ages of C57BL/6 mice receiving injection of the recombinant plasmid of the present invention also show similar results of age-dependent and carrier rate relationship to that of human exposed to HBV under natural infection condition. Therefore, it is preferred to use young C57BL/6 mice in construction of the mouse model for the persistent expression of HBV antigens in the present invention. However, it is impossible to perform tail vein hydrodynamic injection in immature mice, thus 6- to 8- weeks old mice are preferred choice of the present invention.

By analysis of the HBsAg and HBV DNA produced in the bloodstream during HBV virus particles replication process, we prove that persistent expression of HBV antigens for more than 6 months in the transgenic mice of the present invention is associated with the transfection of the chimeric plasmid of the present invention, hence the transgenic animal model of the present invention can be used to elucidate chronic hepatitis mechanism, drug discovery as well as drug evaluation.

Because the present invention is based on the animal model with healthy immunity, thus the induced liver histological and

5

serological status is similar to that of healthy HBV carrier. Consequently the present invention is an ideal model for mechanistic chronic hepatitis studies of hepatitis mechanism and drug evaluation.

EXAMPLE 1

Construction of Vector

The HBV 1.2-full-length DNA is isolated from the plasmid pHBV-48, containing a greater-than-genome-length HBV fragment (subtype adw, gene type A) cloned in pGEM-3Z vector. The replication-competent HBV 1.2-full-length genome replaced the GFP fragment in AAV-GFP vector by ligation of the BamHI/EcoRI (1.8 kb) and EcoRI/BglII (2.0 kb) fragments of pHBV-48 to the BglII site of AAV-GFP vector. The resulting pAAV/HBV (1400) contains the HBV fragment spanning from nucleotides 1400-3182/1-1987 flanked by inverted terminal repeats (ITRs) of AAV.

EXAMPLE 2

Transfection Procedure and Determination

In vivo experiment of the vector transfection, 6 to 8 weeks old male C57BL/6 mice with H-2^b haplotype are used as experimental group, whereas 6 to 8 weeks old male BALB/c with H-2^d haplotype are selected as control group. Ten micrograms of the abovementioned HBV plasmid pAAV/HBV (1400) were injected into the tail vein of mice in a volume of phosphate-buffered saline (PBS) equivalent to 8% of the mouse body weight under hydrodynamic condition that is well known for those skilled in the art. The total volume is delivered within 5 seconds.

After injection, the mice are regularly bled to follow serum concentration of HBV surface antigen (HBsAg), anti-HBV core protein antibodies (anti-HBc) and anti-HBs antibodies (anti-HBs) at the indicated time points, using the AXSYM system kit (Abbott, GmbH Diagnostica). The livers of mice are preserved in optimal cutting temperature (OCT) for immunohistochemical analysis. Expression of HBcAg and HBsAg in liver of the injected mice is visualized by immunohistochemical staining of liver tissue embedded in OCT by using rabbit anti-HBc or anti-HBs antibodies and Envision_{System} HRP (DAB), DAKO Corp., Carpinteria, Calif.).

Changes of serum concentration of HBsAg in experimental group of C57BL/6 mice and the control group BALB/c mice after receiving hydrodynamic injection of the recombinant plasmid of the present invention are shown in FIG. 1. As shown in FIG. 1, when BALB/c (H-2^d) received pAAV/HBV (1400) injection, their HBsAg level increased promptly within one week but dropped quickly in two to three weeks. All the mice developed anti-HBs within 14 days.

When the same plasmid of the present invention is injected into experimental group of C57BL/6 mice, their HBsAg level declined much slower. Notably, in about 40% of recipients, the HBsAg could persist for more than 6 months, meeting the definition of persist human HBV infection. In addition, all C57BL/6 transgenic mice receiving hydrodynamic injection of pAAV/HBV plasmid do not develop anti-HBsAb until day 28 after hydrodynamic injection, despite all of them develop anti-HBV core Ab as BALB/c mice.

EXAMPLE 3

The HBV expression cassette is excised by SmaI digestion (located inside of the two ITRs) from the pAAV/HBV (1400)

6

and subcloned into SmaI site of pGEM4Z to result in pGEM4Z/HBV (1400). Then the obtained recombinant plasmid pGEM4Z/HBV (1400) is injected into the tail vein of C57BL/6 mice according to the method described in example 2.

The results show that the injection of pGEM4Z/HBV (1400) into C57BL/6 mice produced only transient surface antigenemia (shown in FIG. 1b) and all of them developed anti-HBs within 4 weeks (Table 1), suggesting that persistence of HBV in C57BL/6 mice is associated with of injection of pAAV/HBV (1400) plasmid.

EXAMPLE 4

HBV viral RNA transcripts and replication intermediates in the liver of C57BL/6 mice receiving injection of recombinant plasmid pAAV/HBV (1400) are detected by analysis of total liver RNA and DNA using well-known Northern Blotting and Southern blotting. In addition, HBsAg and HBcAg in the hepatocytes are examined by immunohistochemical staining. The results are shown in FIG. 2 and FIG. 3.

As shown in FIG. 2, HBV replication intermediate (including single stranded DNA, double-stranded circular DNA etc.) and RNA transcripts (including 2.1 Kb, 2.4 kb, and 3.5 kb RNA fragment) are detected in the liver of HBsAg-positive C57BL/6 mice at day 70, 93 and 222 after hydrodynamic injection. In addition, HBsAg and HBcAg are also detected in such carrier mice at day 381 after injection (FIG. 3). The results indicate that HBV persistently replicated and survived in the host hepatocytes. Furthermore, long-term expression of HBV in these mice seemed not to cause liver damage as shown by normal ALT level and minimal liver histological change (data not shown).

EXAMPLE 5

According to the method described in example 2, 6.5-week-old male C57BL/6 mice and 13 week-old male C57BL/6 mice, respectively, receive hydrodynamic injection of pAAV/HBV (1400) plasmid in tail vein. Then changes of serum concentration of HBsAg are monitored and analyzed. The results are shown in FIG. 4.

As shown in FIG. 4, equal amount of HBsAg are detected in the bloodstream of both 6.5- and 13-week-old male C57BL/6 mice in the first week after hydrodynamic injection of pAAV/HBV (1400) plasmid. The HBsAg drops quickly to undetectable level in the group of 13-week-old male mice and more than 90% of them developed neutralizing antibodies (anti-HBs) within 56 days after hydrodynamic injection (FIG. 4b). Only 30% of the 6.5-week-old male mice clear HBsAg in their bloodstream and these mice are also anti-HBs positive (FIG. 4 and Table 2).

EXAMPLE 6

6.5-week-old male and female C57BL/6 mice are hydrodynamically injected with pAAV/HBV (1400) plasmid, respectively, according to the method described in example 2. Then changes of serum concentration of HBsAg are monitored and analyzed. The results are shown in FIG. 5.

As shown in FIG. 5, after hydrodynamic injection of pAAV/HBV (1400) plasmid, female C57BL/6 mice have lower level of HBsAg than that of male mice. Furthermore, comparing to male mice, female C57BL/6 mice clear HBsAg more rapidly. More than 40% of male C57BL/6 mice are still HBsAg-positive, while less than 10% of the female C57BL/6 mice show HBsAg-positive after hydrodynamic injection.

TABLE 1

The kinetics of antibody production of C57BL/6 and BALB/c mice

Strain	Plasmid	Anti-HBc Antibody		Anti-HBs Antibody		
		Day 1	Day 7	Day 3	Day 14	Day 28
C57BL/6	pAAV/ HBV(1400)	0/9	9/9	0/9	0/9	0/9
C57BL/6	pGEM4Z/ HBV(1400)	0/4	4/4	0/4	0/4	4/4
BALB/c	pAAV/ HBV(1400)	0/9	9/9	0/9	9/9	9/9

TABLE 2

The kinetics of antibody production of C57BL/6 with different age

Age	Anti-HBc Antibody		Anti-HBs Antibody				
	Day 1	Day 7	Day 3	14	Day 28	Day 42	Day 56
6.5 wks	0/14	14/14	0/14	0/14	1/14	4/14	5/14
13 wks	0/9	9/9	0/9	0/9	3/9	8/9	9/9

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 1

<210> SEQ ID NO 1

<211> LENGTH: 8058

<212> TYPE: DNA

<213> ORGANISM: hepatitis B virus

<400> SEQUENCE: 1

```

aattcccatc atcaataata taccttattt tggattgaag ccaatatgat aatgaggggg 60
tggagtttgt gacgtggcgc ggggcgtggg aacggggcgg gtgacgtagt agtctctaga 120
ggteccacgc gaccttgacg ggcacatgcc cggcatttct gacagctttg tgaactgggt 180
ggccgagaag gaatgggagt tgccgccaga ttctgacatg gatctgaatc tgattgagca 240
ggcaccctcg accgtggcgc agaagctgca tcgctggcgt aatagcgaag aggccccgac 300
cgatcgccct tcccaacagt tgcgcagcct gaatggcgaa tgggaattcca gacgattgag 360
cgtcaaaatg taggtatttc catgagcgtt tttcctgttg caatggctgg cggtaaatatt 420
gttctggata ttaccagcaa ggcgatagtg ttgagttcct ctactcaggc aagtgatgtt 480
attactaatc aaagaagtat tgcgacaacg gtttaattgc gtgatggaca gactctttta 540
ctcgtgggcc tcaactgatta taaaaacact tctcaggatt ctggcgtagc gttcctgtct 600
aaaatccctt taatggcct cctgttttagc tcccgetctg attctaacga ggaaagcagc 660
ttatacgtgc tcgtcaaaagc aaccatagta cgcgcctctg agcggcgcac taagcgcggc 720
gggtgtggtg gttacgcgca gcgtgaccgc tacacttgcc agcgccttag cgcgccctcc 780
tttegttttc ttcccttctt ttctcgccac gttcgcggc tttecccgtc aagctctaaa 840
tcgggggctc cctttagggt tccgatttag tgetttacgg cacctcgacc ccaaaaaact 900
tgattagggt gatggttccac gtagtgggcc atcgccctga tagacggttt ttcgcccttt 960
gacgttgtag tccacgttct ttaatagtg actctgttc caaactggaa caaactcaa 1020
ccctatctcg gtctattctt ttgatttata agggattttg ccgatttcgg cctattgggt 1080
aaaaaatgag ctgatttaac aaaaatttaa cgcgaatttt aacaaaatat taacgtttac 1140
aatttaataa tttgcttata caatcttctt gtttttgggg cttttctgat tatcaaccgg 1200
ggtacatatg attgacatgc tagttttacg attaccgttc atcgccctgca gggggggggg 1260
gggggggggt ggccactccc tctctgccc ctegetoget caactgagcc gggcgaccaa 1320
aggtcgcccg acgcccgggc tttgcccggg cggcctcagt gagcgagcga gcgcgcagag 1380
agggagtggc caactccatc actaggggtt cctagatcct tcgcccggacg tcctttgttt 1440

```

-continued

acgtcccgtc	ggcgctgaat	cccgcggacg	accctctctg	gggcccgttg	ggactctctc	1500
gtccctctct	ccgtctgccg	ttccagccga	ccacggggcg	cacctctctt	tacggggtct	1560
cccgtctgt	gccttctcat	ctgccggtcc	gtgtgcactt	cgcttcacct	ctgcacgttg	1620
catggagacc	accgtgaacg	cccacagat	cctgcccagg	gtcttacata	agaggactct	1680
tggactccca	gcaatgtcaa	cgaccgacct	tgaggcctac	ttcaaagact	gtgtgtttaa	1740
ggactgggag	gagctggggg	aggagattag	gttaaaggtc	tttgtattag	gaggctgtag	1800
gcataaattg	gtctgcgcac	cagcaccatg	caactttttc	acctctgcct	aatcatctct	1860
tgtacatgtc	ccactgttca	agcctccaag	ctgtgccttg	gggtggcttg	gggcatggac	1920
attaccctta	taaagaattt	ggagctaactg	tggagttact	ctcgtttttg	ccttctgact	1980
ttttctcttc	cgtcagagat	ctcctagaca	ccgcctcagc	tctgtatgag	aagccttaga	2040
gtctcctgag	cattgctcac	ctcaccatac	tgactcagg	caagccattc	tctgctgggg	2100
ggaattgatg	actctagcta	cctgggtggg	taataatttg	gaagatccag	catctagggg	2160
tcttgtagta	aattatgtta	atactaactg	gggtttaaag	atcaggcaac	tattgtgggt	2220
tcatatatct	tgcttactt	ttggaagaga	gactgtactt	gaatatttgg	tctctttcgg	2280
agtgtggatt	cgcaactcct	cagcctatag	accaccaaat	gcccctatct	tatcaacact	2340
tccggaaact	actgttggtta	gacgacggga	ccgaggcagg	tcccctagaa	gaagaactcc	2400
ctcgcctcgc	agacgcagat	ctcaatcgcc	cggtcgcaga	agatctcaat	ctcgggaatc	2460
tcaatgttag	tattccttgg	actcataagg	tgggaaactt	tacggggctt	tattcttcta	2520
cagtacctat	ctttaactct	gaatggcaaa	ctccttctct	tcctaagatt	catttacaag	2580
aggacattat	taatagtggt	caacaatttg	tgggccctct	cactgtaaat	gaaaaaagaa	2640
gattgaaatt	aattatgcct	gctagattct	atcctacceca	cactaaatat	ttacccttag	2700
aaaagggaat	taaaccttat	tatccagatc	aggtagttaa	tcattacttc	caaaccagac	2760
attatttaca	tactctttgg	aaggctggta	ttctatataa	gagggaaacc	acacgtagcg	2820
catcattttg	cggttcacca	tattcttggg	aacaagagct	acagcatggg	aggttgggtca	2880
tccaaaacct	gcaaaggcat	ggggacgaat	ctttctgttc	ccaaccctct	gggattcttt	2940
cccgatcacc	agttggacc	tgcatcggga	gccaactcaa	acaatccaga	ttgggacttc	3000
aaccocatca	agaaccactg	gccagcagcc	aaccaggtag	gagtgggagc	attcgggcca	3060
ggactcacc	ctccacacgg	cggtattttg	gggtggagcc	ctcatgctca	gggcatattg	3120
accacagtgt	caacaattcc	tcctctctgc	tccaccaatc	ggcagtcagg	aaggcagcct	3180
actccatct	ctccacctct	aagagacagt	catcctcagg	ccatgcagtg	gaattccact	3240
gccttcaccc	aagctctgca	ggatccocaga	gtcaggggtc	tgtatcttcc	tgtgtgtggc	3300
tccagttcag	gaacagtaaa	ccctgctccg	aatattgcct	ctcacatctc	gtcaatctcc	3360
gcgaggactg	gggaccctgt	ggcgaacatg	gagaacatca	catcaggatt	cctaggaccc	3420
ctgctcgtgt	tacagggggg	gtttttcttg	ttgacaagaa	tcctcacaat	accgcagagt	3480
ctagactcgt	ggtggacttc	tctcaatttt	ctagggggat	caccctgtgt	tcttggtccaa	3540
aattcgcagt	ccccaacctc	caatcactca	ccaacctctc	gtcctccaat	ttgtcctggt	3600
tatcgttggg	tgtgtctgcg	gcgttttctc	atattctctc	tcatcctgct	gctatgcttc	3660
atcttcttat	tggttcttct	ggattatcaa	ggatgtttgc	ccgtttgtcc	tctaattcca	3720
ggatcaacaa	caaccagtac	gggaccatgc	aaaacctgca	cgactcctgc	tcaaggcaac	3780

-continued

tctatgttcc	cctcatgttg	ctgtacaaaa	cctacggatg	gaaattgcac	ctgtattccc	3840
atcccatcgt	cctgggcttt	cgcaaaatc	ctatgggagt	gggcctcagt	ccgtttctct	3900
tggctcagtt	tactagtgcc	atctgttcag	tggttcgtag	ggctttcccc	cactgtttgg	3960
ctttcagcta	tatggatgat	gtggtattgg	gggccaagtc	tgtacagcat	cgtgagtccc	4020
tttataccgc	tgttaccaat	tttcttatgt	ctctgggtat	acatttaaac	cctaacaaaa	4080
caaaaagatg	gggttattcc	ctaaaactca	tgggttacct	aattggaagt	tggagaactt	4140
tgccacagga	tcatattgta	caaaagatca	aacactgttt	tagaaaactt	cctgttaaca	4200
ggcctattga	ttggaagta	tgtcaaagaa	ttgtgggtct	tttgggcttt	gctgctccat	4260
ttacacaatg	tggatctcct	gccttgatgc	ctttgatgc	atgtatacaa	gctaaacagg	4320
ctttcacttt	ctcgccaact	tacaaggcct	ttctaagtaa	acagtacatg	aacctttacc	4380
ccgttgctcg	gaaacggcct	ggtctgtgcc	aagtgtttgc	tgacgcaacc	cccactggct	4440
ggggcttggc	catagcccat	cagcgcacgc	gtggaacctt	tgtggctcct	ctgccgatcc	4500
atactgcgga	actcctagcc	gcttgttttg	ctcgcagccg	gtctggggca	aagctcatcg	4560
gaactgacaa	ttctgtcgtc	ctctcgcgga	aatatacatc	atttccatgg	ctgctaggct	4620
gtactgccaa	ctggatcctt	cgcgggacgt	cctttgttta	cgtcccgtcg	gcgctgaatc	4680
ccgcggaaga	cccctctcgg	ggcgccttgg	gactctctcg	tcccctctc	cgtctgccgt	4740
tccagccgac	cacggggcgc	acctctcttt	acgcgggtctc	cccgtctgtg	ccttctcacc	4800
tgcgggtccg	tgtgcacttc	gcttcacctc	tgcacgttgc	atggagaacca	ccgtgaacgc	4860
ccatcagatc	ctgcccgaag	tcttacataa	gaggactcct	ggactcccag	caatgtcaac	4920
gaccgacctt	gaggcctact	tcaaagactg	tgtgtttaag	gactgggagg	agctggggga	4980
ggagattagg	ttaaaggctc	ttgtattagg	aggctgtagg	cataaattgg	tctgcgcacc	5040
agcaccatgc	aactttttca	cctctgccta	atcatctctt	gtacatgtcc	cactgttcaa	5100
gcctccaage	tgtgccttgg	gtggctttgg	ggcatggaca	ttgacctta	taaagaattt	5160
ggagctactg	tggagtactc	ctcgtttttg	ccttctgact	ttttccttc	cgtcagagat	5220
ctaggaaccc	ctagtgatgg	agttggccac	tccctctctg	cgcgctcgct	cgtcactga	5280
ggcgcgccgg	gcaaagcccg	ggcgtcgggc	gacctttggt	cgcccggcct	cagtgaacga	5340
gcgagcgcgc	agagaggggg	tggccaaccc	cccccccccc	cccctgcag	gcgattctct	5400
tgtttgtctc	agactctcag	gcaatgacct	gatagccttt	gtagagacct	ctcaaaaaa	5460
gctaccctct	ccggcatgaa	tttatcagct	agaacggttg	aatatcatat	tgatggtgat	5520
ttgactgtct	ccggccttcc	tcaccogttt	gaatctttac	ctacacatta	ctcaggcatt	5580
gcatttaaaa	tatatgaggg	ttctaaaaat	ttttatcctt	gcgttgaaat	aaaggcttct	5640
cccgcaaaag	tattacaggg	tcataatggt	tttggtacaa	ccgatttagc	tttatgctct	5700
gaggctttat	tgettaattt	tgetaattct	ttgccttgcc	tgtatgattt	attggatggt	5760
ggaattcctg	atgcggatatt	ttctccttac	gcactctgtgc	ggatattcac	accgcatatg	5820
gtgcaactctc	agtacaactct	gctctgatgc	cgcatagtta	agccagcccc	gacaccgcgc	5880
aacaccgcct	gacgcgcctc	gacgggcttg	tctgctcccg	gcctccgctt	acagacaagc	5940
tgtgaccgtc	tccggagctc	gcattgtgtca	gaggttttca	ccgtcatcac	cgaacgcgc	6000
gagacgaaag	ggcctcgtga	tacgcctatt	tttatagggt	aatgtcatga	taataatggt	6060

-continued

ttcttagacg tcaggtggca cttttcgggg aaatgtgcgc ggaaccoccta tttgtttatt	6120
tttctaaata cattcaaata tgtatccgct catgagacaa taaccctgat aaatgcttca	6180
ataatattga aaaaggaaga gtatgagat tcaacatttc cgtgtcgccc ttattccctt	6240
ttttgcggca ttttgccctc ctgtttttgc tcaccagaa acgctggtga aagtaaaaga	6300
tgctgaagat cagttgggtg cacgagtggg ttacatcgaa ctggatctca acagcggtaa	6360
gatccttgag agttttcgcc ccgaagaacg ttttccaatg atgagcactt ttaaagtctt	6420
gctatgtggc gcggtattat cccgtattga cgccgggcaa gagcaactcg gtcgccgat	6480
acactattct cagaatgact tggttgagta ctcaccagtc acagaaaagc atcttacgga	6540
tggcatgaca gtaagagaat tatgcagtgc tgccataacc atgagtgata acactgcggc	6600
caacttactt ctgacaacga tcggaggacc gaaggagcta accgcttttt tgcacaacat	6660
gggggatcat gtaactcgcc ttgatcgttg ggaaccggag ctgaatgaag ccataccaaa	6720
cgacgagcgt gacaccacga tgcctgtagc aatggcaaca acgttgcgca aactattaac	6780
tggcgaacta ctactctag cttcccgga acaattaata gactggatgg aggcggataa	6840
agttgcagga ccacttctgc gctcggccct tccggctggc tggtttattg ctgataaatc	6900
tggagccggt gagcgtgggt ctgcgggtat cattgcagca ctggggccag atggttaagc	6960
ctcccgatc gtagttatct acacgacggg gagtcaggca actatggatg aacgaaatag	7020
acagatcgct gagataggtg cctcactgat taagcattgg taactgtcag accaagtta	7080
ctcatatata ctttagattg atttaaaact tcatttttaa tttaaaagga tctaggtgaa	7140
gatccttttt gataatctca tgacaaaaat cccttaacgt gagtttctgt tccactgagc	7200
gtcagacccc gtagaaaaga tcaaaggatc ttcttgagat cctttttttc tgcgcgtaat	7260
ctgctgcttg caaacaaaa aaccaccgct accagcggtg gtttgtttgc cggatcaaga	7320
gctaccaact ctttttccga aggtaactgg cttcagcaga gcgcagatac caaatactgt	7380
ccttctagtg tagccgtagt taggccacca cttcaagaac tctgtagcac cgctacata	7440
cctcgctctg ctaatcctgt taccagtggc tgctgccagt ggcgataagt cgtgtcttac	7500
cgggttggac tcaagacgat agttaccgga taaggcgcag cggcgggct gaaacggggg	7560
ttcgtgcaca cagcccagct tggagcgaac gacctacacc gaactgagat acctacagcg	7620
tgagctatga gaaagcgcca cgcttccga agggagaaag gcggacaggt atccggttaag	7680
cggcagggtc ggaacaggag agcgcacgag ggagcttcca ggggaaacg cctggtatct	7740
ttatagtcct gtcgggttcc gccacctctg acttgagcgt cgatttttgt gatgctctc	7800
agggggcgcg agcctatgga aaaacgccag caacgcggcc tttttacggt tectggcctt	7860
ttgtggcct tttgctcaca tgttctttcc tgcgttatcc cctgattctg tggataaccg	7920
tattaccgcc tttgagttag ctgataccgc tcgccgcagc cgaacgaccg agcgcagcga	7980
gtcagtgagc gaggaagcgg aagagcgccc aatacgcaaa ccgectctcc ccgcgcttg	8040
gccgattcat taatgcag	8058

What is claimed is:

1. A method for expressing hepatitis B viral antigen in vivo, comprising the steps of:

- (1) providing a recombinant plasmid having the sequence of SEQ ID NO: 1; and
- (2) delivering the recombinant plasmid into a male C57BL/6 mouse and transfecting the hepatocytes of the mouse by hydrodynamic injection;

60

wherein an expression of hepatitis B viral antigens persists for more than 26 weeks in the hepatocytes.

2. The method as claimed in claim 1, wherein the mouse is H-2^b haplotype.

65

3. The method as claimed in claim 1, wherein the age of the mouse is aged 6 to 8 weeks.

15

4. The method as claimed in claim 1, wherein the recombinant plasmid is prepared in a pre-mix solution before delivering into the mouse.

5. The method as claimed in claim 4, wherein the solution is phosphate buffer.

6. The method as claimed in claim 1, wherein the recombinant plasmid is injected into a tail vein of the mouse.

16

7. The method as claimed in claim 1, wherein the hepatitis B viral antigen is hepatitis B viral surface antigen.

8. The method as claimed in claim 1, wherein the hepatitis B viral antigen is hepatitis B viral core protein.

* * * * *